



# IX Congresso Ibérico das Ciências do Solo

**“O solo, recurso estratégico  
para uma sociedade sustentável”**

**22 a 24 de junho de 2022**

Oeiras - Lisboa

# Comunicações

## **Ficha Técnica**

### **Edição**

Sociedade Portuguesa da Ciência do Solo

### **Título**

Comunicações do IX Congresso Ibérico das Ciências do Solo  
O solo, recurso estratégico para uma sociedade sustentável

### **Capa e paginação**

Rui Tomás Monteiro

©

### **Comissão Organizadora**

Adélia Varela (INIAV); Ana Marta Paz (SPCS, INIAV); Carlos Alexandre (SPCS, UÉvora); Clara Pinto (INIAV); Erika Santos (ISA - ULisboa); Jorge Mataix Solera (SECS, UMH); Jorge Nunes (ISA - ULisboa); Maria do Carmo Horta (SPCS, IPCastelo Branco); Maria Madalena Fonseca (ISA - ULisboa); Nádia Castanheira (INIAV); Nuno Cortez (SPCS, ISA - ULisboa); Paula Alvarenga (SPCS, ISA - ULisboa)

### **Comissão Científica**

Amaia Nogales (ISA- ULisboa); Ana Marta Paz (INIAV); Carlos Alexandre (UÉvora); David Badía Villas (SECS - UZaragoza); David Fangueiro (ISA - ULisboa); Engracia Madejón Rodríguez (SECS, IRNAS-CSIC, Sevilla); Erika Santos (ISA - ULisboa); Ernesto Vasconcelos (ISA - ULisboa); Gael Bárcenas Moreno (SECS, USevilla); Gonçalo Rodrigues (ISA - ULisboa); Henrique Ribeiro (ISA - ULisboa); Irene Ortiz Bernad (SECS, UGranada); Isabel Maria Oliveira Brito (UÉvora); João Coutinho Mendes (UTAD); Jorge Mataix-Solera (SECS, UMiguel Hernández); Jorge Pinheiro (UAçores); José Alexandre Andrade (UÉvora); José Casimiro Martins (INIAV); José Manuel Rato Nunes (IP Portalegre); Manuel Jordán Vidal (UMiguel Hernández); Manuel Madeira (ISA - ULisboa); Maria da Conceição Gonçalves (INIAV); Maria do Carmo Horta (IP Castelo Branco); Maria Madalena Fonseca (ISA - ULisboa); Maria Manuela Abreu (ISA - ULisboa); Miguel Brito (IP Viana do Castelo); Minerva García-Carmona (UMiguel Hernández); Nuno Cortez (ISA - ULisboa); Oriol Ortiz Perpiñá (SECS, UZaragoza); Paula Alvarenga (ISA - ULisboa); Ruth Pereira (UPorto); Sara Ibáñez Asensio (SECS, UPValència); Selma B. Pena (ISA - ULisboa); Tiago Ramos (MARE-TEC, IST - ULisboa); Tomás Figueiredo (IP Bragança)



Dia 24/06/2022

## SESSÃO 2 – Sala 2

### Tema 4 (T4) - Solos e temas ambientais / Problemas ambientales y de suelos

Sala	Nº	Tema	Título	Apresentado por
2	P17	T4	Extractabilidad de metales pesados en suelos urbanos	María Roo Calvo
2	P18	T4	Genes de resistencia a antibióticos en los suelos: ¿Es posible su transferencia a las plantas?	Maria Teresa Gomez Sagasti
2	P19	T4	Evaluación del contenido de glomalina y su relación con la agregación en suelos urbanos	Remigio Paradelo Núñez
2	P20	T4	Short-term evolution of physico-chemical properties of Technosols made from contaminated soils by pyritic sludge	Antonio Aguilar Garrido
2	P21	T4	Restauración de estériles de mina con enmiendas orgánicas y especies vegetales: Impacto sobre las propiedades físicas del suelo	Carmela Monterroso
2	P22	T4	Remediation of a chlorpyrifos contaminated soil using novel bacterial strains and cyclodextrin. Evaluation of its effectiveness by ecotoxicity studies	Alba Lara-Moreno
2	P23	T4	Red de estaciones de medida de temperatura y humedad del suelo	Dolores Boquera Ferreiro
2	P24	T4	Os microrganismos do solo no montado: uma estratégia para adaptação às alterações climáticas	Márcia de Castro Silva
2	P25	T4	La fitorremediación de suelos afectados por la actividad cinegética: lecciones aprendidas, retos y futuras oportunidades	Maria Teresa Gomez Sagasti
2	P26	T4	Biorremediación de suelos contaminados con aceite de motor mediante bioestimulantes obtenidos de lodos de depuradora	Isidoro A. Gomez Parrales
2	P27	T4	Comportamiento de oxifluorfen en suelos enmendados con diferentes fuentes de materia orgánica y sometido a sequía severa	Isidoro A. Gomez Parrales
2	P28	T4	Soil and vegetation recovery in an area affected by residual pollution after remediation measures	Mario Paniagua López
2	P29	T4	Utilização de parâmetros físico-químicos e ecotoxicológicos na avaliação da remediação de solos degradados por atividades mineiras	Adriana Catarino
2	P30	T4	Cu e Zn em frações físicas de solos de vinhedos no Sul do Brasil	Cledimar Rogério Lourenzi
2	P31	T4	Toxicidad de amoxicilina sobre las comunidades bacterianas del suelo en suelos agrícolas	Vanessa Santás Miguel
2	P32	T4	Disipación asistida de un mix de PAHs en suelos contaminados. Efecto del tipo de suelo y compuestos que incrementan su disponibilidad	Fernando Madrid Díaz
2	P33	T4	Nematodes as soil status bioindicators: MaisSolo project	Maria Lurdes Inácio

## Soil and vegetation recovery in an area affected by residual pollution after remediation measures

### *Recuperación del suelo y la vegetación en una zona afectada por contaminación residual tras la aplicación de medidas de remediación*

**Paniagua-López, Mario<sup>1\*</sup>; García-Robles, Helena<sup>2</sup>; Aguilar-Garrido, Antonio<sup>1</sup>; Romero-Freire, Ana<sup>1</sup>; Lorite, Juan<sup>2</sup>; Sierra-Aragón, Manuel<sup>1</sup>**

<sup>1</sup> Departamento de Edafología y Química Agrícola, Universidad de Granada, Campus Fuentenueva s/n, 18071 Granada, Spain.

<sup>2</sup> Departamento de Botánica, Universidad de Granada, Campus Fuentenueva s/n, 18071 Granada, Spain. \*mpaniagua@ugr.es

#### **Abstract**

The present study evaluates the effectiveness of remediation treatments applied to residual polluted soils affected by the Aznalcóllar mining spill occurred in 1998 (SW Spain). For this purpose, the influence of the treatments on the main soil properties and on the evolution of natural vegetation was studied. The results showed that amendments application improved the main soil properties and reduced soil toxicity, which led to a partial natural recovery of the vegetation cover and species richness in amended soils. *Spergularia rubra* and *Lamarckia aurea*, two of the main plant species present in the area and more strongly associated to high pollutant concentrations, showed a high bioaccumulation factor (BAF) for the main pollutants in the area (Pb and As) in shoots and, specially, in roots. *Lamarckia aurea* showed a remarkably high accumulation of these elements in roots, which is consistent with its distribution, strongly associated to areas where the vegetation growth is more restricted and where no other plant species are able to grow. Thus, both *Spergularia rubra* and *Lamarckia aurea* are important key species in this area, which may favour the subsequent colonization of the residual polluted soils by other less tolerant species to the pollutants present in these soils.

Keywords: Soil pollution, amendments, natural remediation, *Lamarckia aurea*, bioaccumulation factor

#### **Resumen**

El presente estudio evalúa la eficacia de diferentes tratamientos de remediación aplicados en suelos residualmente contaminados afectados por el vertido minero de Aznalcóllar, ocurrido en 1998 (SO de España). Se estudió la influencia de estos tratamientos sobre las principales propiedades del suelo y sobre la evolución de la vegetación. Los resultados mostraron que las enmiendas mejoraron las principales propiedades del suelo y redujeron su toxicidad, lo que condujo a una recuperación natural parcial de la cubierta vegetal y la riqueza de especies en los suelos enmendados. *Spergularia rubra* y *Lamarckia aurea*, dos de las principales especies vegetales presentes en la zona y más fuertemente asociadas a altas concentraciones de contaminantes, mostraron un elevado factor de bioacumulación (BAF) para los principales contaminantes en la zona (Pb y As) tanto en parte aérea como en raíz. *Lamarckia aurea* mostró una elevada acumulación de estos elementos en las raíces, lo que concuerda con su distribución, fuertemente asociada a zonas donde el crecimiento de la vegetación está más limitado. *Spergularia rubra* y *Lamarckia aurea*, por tanto, son especies clave en esta zona, ya que pueden favorecer la colonización posterior de estos suelos residualmente contaminados por parte de otras especies menos tolerantes a la contaminación.

Palabras clave: Contaminación del suelo, enmiendas, remediación natural, *Lamarckia aurea*, factor de bioacumulación

## Introduction

Soil pollution by heavy metals is a worldwide problem which poses a risk to ecological systems and to human health and food production. Anthropogenic activities, such as mining, represent important potential sources of soil pollution (Dermont et al., 2008; Liu et al., 2018). One of the most important mining accidents worldwide associated with metal mining spill occurred in Aznalcóllar (SW Spain) on 25 April 1998, when the tailing dam collapsed and around  $4.5 \times 10^6 \text{ m}^3$  of acidic waters and toxic tailings were spilled into the Agrio and Guadiamar river basins, affecting an area of over  $43 \text{ km}^2$  (Simón et al., 1999).

A great variety of soil remediation techniques, both in-situ and ex-situ, have been developed and tested in contaminated soils with the aim of reducing the polluting effects (Liu et al., 2018). These techniques may be based on the addition of organic and inorganic amendments to assist the natural remediation processes of soils damaged or contaminated by pollutants (Adriano et al., 2004).

Plants, as an element of the ecosystem, play a significant role in the restoration of degraded or polluted areas, and may act as an indicator of the degree of contamination present in soils. They can accumulate trace elements, especially heavy metals, in or on their tissues, and have a great ability to adapt to variable degrees of pollution present in the environment (Kabata-Pendias, 2000). Also, passive restoration by native vegetation that spontaneously colonize a contaminated area, in improving conditions, is an important factor to consider when evaluating the success of remediation actions (Álvarez-Rogel et al., 2021), as they may act as nurse plants and thus facilitate the growth of less stress-tolerant species (Navarro-Cano et al., 2018).

The present study aims to evaluate the effectiveness of remediation treatments applied to residual polluted soils affected by the Aznalcóllar mining spill by analyzing their influence on the main soil properties, as well as in the spontaneous vegetation recovery.

## Material and methods

The study site corresponds to the area affected by the toxic spill of the Aznalcóllar mine (SW Spain). The area nearby the tailing dam where the spill occurred is characterised by the existence of numerous randomly distributed unvegetated patches. For this study, three unvegetated patches of soils which remained affected by residual pollution were selected. In each of them, an experimental plot of  $24 \text{ m}^2$  was established, subdivided into 6 subplots where six different treatments were applied. The treatments consisted of: BS: Biopile, which consisted of a mixture of contaminated soil affected by the spill with the adjacent recovered soil in a proportion of 50% (w/w); BVS: Biopile + vermicompost; GS: Gypsum mining spoil; GVS: Gypsum mining spoil + vermicompost; MS: Marble sludge; MVS: Marble sludge + vermicompost.

Eighteen months after the application of the treatments, composite samples of the topsoil (0 – 10 cm) for each of the treatments were taken, and the main soil properties were analyzed. Also, the vegetation present in the plots was monitored, by measuring the vegetation cover and the plant species richness on every treatment applied. For this purpose, a  $0.25 \text{ m}^2$  grid divided into 100 cells was used, and the number of cells occupied by herbaceous species, both as total cover and as cover per species, as well as the total number of different species present on the grid, were counted. This was replicated three times per experimental plot and treatment. Finally, vegetation samples of the most frequent plant species were collected to analyze the pollutant elements uptake by these colonizing species. Bioaccumulation factor (BAF), both for aerial part and for roots, was calculated to determine the plant's capacity to take up metal from the soil and to either transport it to the aboveground parts or to accumulate it in the roots. To do so, the different parts of the plant samples were digested in a microwave XP1500Plus (Mars®) in  $\text{HNO}_3$ :  $\text{H}_2\text{O}_2$  (1:1), and concentrations of potential pollutants (Pb, As, Zn, Cu) were measured by ICP-MS in a PE SCIEX ELAN-5000A spectrometer. The plant-available (EDTA-extractable) metal fraction in soils samples

was also calculated for this end. Finally, translocation factor (TF) was estimated as the ratio between trace element content in aboveground parts of the plants and trace element content in roots, to determine the degree of element migration from the roots to the aerial part of the plant.

## Results and discussion

### Main soil properties related to contamination

Table 1 shows the values of the main soil properties related to the contamination due to the Aznalcóllar mining spill. The soils in which a high degree of contamination persists (CS) were characterized mainly by a strong acid pH and high salinity (EC), as well as by a low organic matter content due to the absence of vegetation cover. Compared to CS, recovered soils (RS) and soils not affected by the spill (US) showed a less acidic pH, but specially a significantly lower salinity and a higher organic matter content. As for the amended soils, pH increased in all cases compared to CS, and reached values even higher than US for treatments with marble (MS and MVS) and gypsum (GS and GVS). EC and organic carbon (OC) content slightly improved under treatments, although OC was higher in soils in which vermicompost was added.

Table 1. Main soil properties (EC: Electrical conductivity; OC: Organic carbon content) related to the Aznalcóllar mining accident.

	pH		EC (dS/m)		OC (%)	
	Mean	sd	Mean	sd	Mean	sd
<b>US</b>	6,06de	0,28	0,08a	0,06	2,32bc	1,60
<b>RS</b>	4,28ab	0,48	0,98b	0,61	3,68c	1,46
<b>CS</b>	3,49a	0,23	3,12d	0,63	0,61a	0,09
<b>BS</b>	5,37cd	0,87	2,35cd	0,30	1,14ab	0,16
<b>BVS</b>	4,96bc	0,41	2,15c	0,51	1,62ab	0,38
<b>GS</b>	6,22def	0,29	2,38cd	0,06	0,89a	0,11
<b>GVS</b>	6,41ef	0,19	2,36cd	0,04	1,67ab	0,26
<b>MS</b>	6,78ef	0,19	2,29c	0,07	1,20ab	0,24
<b>MVS</b>	7,07f	0,07	2,14c	0,26	1,41ab	0,24

US: uncontaminated soil; RS: Recovered soil; CS: Contaminated soil; BS: Biopile soil; BVS: Biopile soil + vermicompost addition; GS: Gypsum amended soil; GSV: GS + vermicompost addition; MS: Marbel amended soil; MVS: MS + vermicompost addition. sd: standard deviation (n=6). Lowercase letters represent significant differences among soils (Kruskal-Wallis, P < 0.05).

### Plant cover and richness and pollutants concentration in plants

Both in US and RS the vegetation cover was total, unlike CS where the vegetation growth was completely inhibited. Species richness compared to US was slightly reduced in RS, while in treated soils it was about half of it. The treatments applied on CS, however, improved soil properties and reduced soil toxicity, allowing vegetation to develop, with a higher percentage of cover and species richness in the vermicompost treatments (Fig. 1).




Fig. 1 - Plant cover and species richness under treatments.

US: uncontaminated soil; RS: Recovered soil; CS: Contaminated soil; BS: Biopile soil; BVS: Biopile soil + vermicompost addition; GS: Gypsum amended soil; GSV: GS + vermicompost addition; MS: Marbel amended soil; MVS: MS + vermicompost addition. Sr: *Spergularia rubra*; La: *Lamarckia aurea*.

*Spergularia rubra* and *Lamarckia aurea* are two species frequently identified in the soils affected by the Aznalcóllar spill associated with areas where pollutant concentrations are high, with acid pH and high salinity. Thus, the capacity of these species as accumulators of pollutants was analyzed. The bioaccumulation factor (BAF) showed that both *S. rubra* and *L. aurea* are potent accumulators of Pb and As, both in the aerial part and, above all, in roots (Table 2). However, for these elements, *L. aurea* showed a higher BAF than *S. rubra*, specially in roots, although the abundance of the latter was higher in all treatments (Fig. 1). This indicates that *L. aurea* is highly tolerant to great concentrations of pollutants, and that the conditions on the amended soils after the treatments has improved, allowing vegetation less tolerant to higher contamination levels to colonize them and to be present at higher rate. This is consistent with previous studies in the

area, that indicated that *L. aurea* has a greater presence in the areas with higher levels of pollution (García-Carmona et al., 2019). For Zn, both species showed BAF values greater than unity, but to a lesser extent than for Pb and As, and none of them showed a significant accumulation capacity for Cu. In addition, the translocation factor (TF) was lower than one in all cases, indicating that there was no migration of these elements from roots to the aboveground parts of these species (Table 2). However, it was observed that TF in *S. rubra* was about twice as high as in *L. aurea* for all elements. This indicates that migration of elements to aerial part in *S. rubra* is higher than in *L. aurea*, where accumulation in roots is greater, which may allow this species to be present in soils with a higher degree of pollution.

Table 2 – Bioaccumulation factor (BAF) and translocation factor (TF) for main plant species present in the study area

		<i>S. rubra</i>		<i>L. aurea</i>	
		mean	sd	mean	sd
<b>BAF</b>	<b>Pb_shoot</b>	32,8	51,2	41,8	55,7
	<b>Pb_root</b>	42,8	42,7	136,7	150,9
<b>TF</b>	<b>Pb</b>	0,6	0,4	0,3	0,2
	<b>As_shoot</b>	14,1	13,6	15	10,9
<b>BAF</b>	<b>As_root</b>	19,1	15,5	39,1	27,9
	<b>TF</b>	<b>As</b>	0,7	0,4	0,4
<b>BAF</b>	<b>Zn_shoot</b>	5,6	4,5	2,9	2,1
	<b>Zn_root</b>	5,0	3,3	4,1	2,2
<b>TF</b>	<b>Zn</b>	1,0	0,5	0,6	0,4
<b>BAF</b>	<b>Cu_shoot</b>	0,8	0,4	0,7	0,5
	<b>Cu_root</b>	0,8	0,4	1,5	0,8
<b>TF</b>	<b>Cu</b>	0,8	0,5	0,4	0,2

## Conclusions

The remediation treatments improved the main soil properties by increasing soil pH value, organic carbon content and reducing salinity. Thus, treatments were also effective in enhancing the spontaneous vegetation recovery of the contaminated soil patches. Vermicompost amendment treatments showed the greatest effectiveness in vegetation recovery, leading to a higher species richness and plant cover in the amended soils. *S. rubra* and *L. aurea* showed a great capacity to accumulate Pb and As both in shoots and roots, which leads them to have a high tolerance to the high concentration of pollutants present in the studied soils. This

points to these species as a key species, as they may act as pioneer species and thus facilitate the secondary succession of the residual polluted soils which remain in the area.

## Acknowledgements

This work was supported by the Research Project RTI 2018-094327-B-I00 (Ministry of Science, Innovation and Universities).

## References

Adriano, D. C., Wenzel, W. W., Vangronsveld, J., & Bolan, N. S. (2004). Role of assisted natural remediation in environmental cleanup. *Geoderma*, 122(2-4), 121-142. <https://doi.org/10.1016/j.geoderma.2004.01.003>

Álvarez-Rogel, J., Peñalver-Alcalá, A., Jiménez-Cárceles, F. J., Tercero, M. C., & González-Alcaraz, M. N. (2021). Evidence supporting the value of spontaneous vegetation for phytomanagement of soil ecosystem functions in abandoned metal(loid) mine tailings. *Catena*, 201, 105191. <https://doi.org/10.1016/j.catena.2021.105191>

Dermont, G., Bergeron, M., Mercier, G., & Richer-Laflèche, M. (2008). Metal-contaminated soils: remediation practices and treatment technologies. *Practice periodical of hazardous, toxic, and radioactive waste management*, 12(3), 188-209. [https://doi.org/10.1061/\(ASCE\)1090-025X\(2008\)12:3\(188\)](https://doi.org/10.1061/(ASCE)1090-025X(2008)12:3(188))

García-Carmona, M., García-Robles, H., Torrano, C. T., Ondoño, E. F., Moreno, J. L., Aragón, M. S., & Peinado, F. M. (2019). Residual pollution and vegetation distribution in amended soils 20 years after a pyrite mine tailings spill (Aznalcóllar, Spain). *Science of the Total Environment*, 650, 933-940. <https://doi.org/10.1016/j.scitotenv.2018.09.092>

Kabata-Pendias, A. (2000). *Trace elements in soils and plants*. Third Edition. CRC Press, Inc. Boca Raton. USA. pp. 365, 413. <https://doi.org/10.1201/9781420039900>

Liu, L., Li, W., Song, W., & Guo, M. (2018). Remediation techniques for heavy metal-contaminated soils: Principles and applicability. *Science of the Total Environment*, 633, 206-219. <https://doi.org/10.1016/j.scitotenv.2018.03.161>

Navarro-Cano, J. A., Verdú, M., & Goberna, M. (2018). Trait-based selection of nurse plants to restore ecosystem functions in mine tailings. *Journal of Applied Ecology*, 55(3), 1195-1206. <https://doi.org/10.1111/1365-2664.13094>

Simón, M., Ortíz, I., García, I., Fernández, E., Fernández, J., Dorronsoro, C., & Aguilar, J. (1999). Pollution of soils by the toxic spill of a pyrite mine (Aznalcóllar, Spain). *Science of the Total Environment*, 242(1-3), 105-115. [https://doi.org/10.1016/S0048-9697\(99\)00378-2](https://doi.org/10.1016/S0048-9697(99)00378-2)

## Apoios

---



Companhia das Lezírias



## Patrocínios

---



aquagri